

Quantum Replicator Dynamics

Esteban Guevara Hidalgo^{†‡}

[†]*Departamento de Física, Escuela Politécnica Nacional, Quito, Ecuador*

[‡]*SIÓN, Autopista General Rumiñahui, Urbanización Edén del Valle,
Sector 5, Calle 1 y Calle A # 79, Quito, Ecuador*

We propose quantization relationships which would let us describe and solution problems originated by conflicting or cooperative behaviors among the members of a system from the point of view of quantum mechanical interactions. The quantum analogue of the replicator dynamics is the equation of evolution of mixed states from quantum statistical mechanics. A system and all its members will cooperate and rearrange its states to improve their present condition. They strive to reach the best possible state for each of them which is also the best possible state for the whole system. This led us to propose a quantum equilibrium in which a system is stable only if it maximizes the welfare of the collective above the welfare of the individual. If it is maximized the welfare of the individual above the welfare of the collective the system gets unstable and eventually it collapses.

PACS numbers: 03.65.-w, 02.50.Le, 03.67.-a, 03.67.Lx

I. INTRODUCTION

Why quantum versions have shown be more efficient and provide more advantages than their classical versions? Maybe quantum mechanics is a more general theory that we had thought and it would let us explain more correctly not only economical but also biological phenomena which have been described before through game theory. If there is an actual connection between game theory and quantum mechanics, we could establish quantization relationships between our classical into quantum systems for any cases. We could also propose a more general quantum equilibrium applicable to any classical system but which remains inside its foundations the classical definitions of equilibrium. The purpose of this paper is show the relationships between the replicator dynamics and quantum mechanics. We will propose the quantization relationships for a classical system and a new quantum equilibrium concept from which is possible to obtain and understand the classical equilibria.

Game theory [1, 2, 3] is the study of decision making of competing agents in some conflict situation. It tries to understand the birth and the development of conflicting or cooperative behaviors among a group of individuals who behave rationally and strategically according to their personal interests. Each member in the group strive to maximize its welfare, state, utilities or payoffs by choosing the best courses of strategies from a cooperative or individual point of view. Game theory has been applied to solve many problems in economics, social sciences, biology and engineering.

Evolutionary game theory [4, 5, 6] does not rely on rational assumptions but on the idea that the Darwinian process of natural selection [7] drives organisms towards the optimization of reproductive success [8]. It combines the principles of game theory, evolution and dynamical systems to explain the distribution of different phenotypes in biological populations. Instead of working out

the optimal strategy, the different phenotypes in a population are associated with the basic strategies that are shaped by trial and error by a process of natural selection or learning. It can also be used to interpret classical games from a different perspective. Instead of directly calculating properties of a game, populations of players using different strategies are simulated and a process similar to natural selection is used to determine how the population evolves. This is made through the stability analysis of differential equations and the implications to the games [9]. The central equilibrium concept of evolutionary game theory is the notion of Evolutionary Stable Strategy introduced by J. Smith and G. Price [4, 10]. An ESS is described as a strategy which has the property that if all the members of a population adopt it, no mutant strategy could invade the population under the influence of natural selection. ESS are interpreted as stable results of processes of natural selection. The natural selection process that determines how populations playing specific strategies evolve is known as the replicator dynamics [5, 6, 9, 11] whose stable fixed points are Nash equilibria [2].

Quantum games have proposed a new point of view for the solution of the classical problems and dilemmas in game theory. It has been shown that quantum games are more efficient than classical games and provide a saturated upper bound for this efficiency [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22]. Meyer [12] quantized a coin tossing game and found out that one player could increase his expected payoff and win with certainty by implementing a quantum strategy against his opponent's classical strategy. Eisert *et al* [13] developed a general protocol for two player-two strategy quantum games with entanglement by quantizing prisoner's dilemma. They found a unique Nash Equilibrium, which is different from the classical one, and the dilemma could be solved if the two players are allowed to use quantum strategies. This also was extended to multi-player games [23]. Marinatto and Weber [15] extended

the concept of a classical two-person static game to the quantum domain by giving an Hilbert structure to the space of classical strategies. They showed that the introduction of entangled strategies in battle of the sexes game leads to a unique solution of this game. Du *et al* [17] implemented a game via nuclear magnetic resonance (NMR) system. It was demonstrated that neither of the two players would win the game if they play rationally, but if they adopt quantum strategies both of them would win. Quantum games has been used to explore unsolved problems of quantum information [24] and in the production of algorithms for quantum computers [25]. Quantum communication can be considered as a game where the objective is maximize effective communication. Also distributed computing, cryptography, watermarking and information hiding tasks can be modeled as games [26, 27, 28, 29, 30, 31, 32]. Piotrowski and Sladkowski have modeled markets, auctions and bargaining assuming traders can use quantum protocols [33, 34]. In the new quantum market games, transactions are described in terms of projective operations acting on Hilbert spaces of strategies of traders. A quantum strategy represents a superposition of trading actions and can achieve outcomes not realizable by classical means [35]. Furthermore, quantum mechanics has features that can be used to model aspects of market behavior. For example, traders observe the actions of other players and adjust their actions [34] and the maximal capital flow at a given price corresponds to entanglement between buyers and sellers [33]. Nature may be playing quantum survival games at the molecular level [36, 37]. It could lead us to describe many of the life processes through quantum mechanics like Gogonea and Merz [38] on protein molecules. Game theory and quantum game theory offer interesting and powerful tools and their results will probably find their applications in computation, complex system analysis and cognition sciences [39, 40, 41].

II. REPLICATOR DYNAMICS & EGT

Payoffs in biological games are in terms of fitness a measure of reproductive success [8]. Strategies are considered to be inherited programs for any conceivable situation which control the individual's behavior. The members of a population interact in game situations and the joint action of mutation and selection replaces strategies by others with a higher reproductive success. It is less important to know which member plays which strategy within a population but it is important to know the relative frequency of actions, it means the probability of playing any strategy.

A *symmetric two - person game* $G = (S, E)$ consists of a finite nonempty pure strategy set S and a payoff function E which assigns a real number to the pair (s_i, s_j) . $E(s_i, s_j)$ is the payoff obtained by a player who plays the strategy s_i against an opponent who plays the strategy

s_j . A *mixed strategy* x is a probability distribution over S .

A *best reply* to q is a strategy p which maximizes $E(p, q)$. An *equilibrium point* is a pair (p, q) with the property that p and q are best replies to each other. A strategy r is a *strict best reply* to a strategy q if it is the only best reply to q . A strict best reply must be a pure strategy. An equilibrium point (p, q) is called strict if p and q are strict best replies to each other. A best reply to p which is different from p is called *alternative best reply* to p .

A *Nash equilibrium* (NE) is a set of strategies, one for each player, such that no player has an incentive to unilaterally change his action. Players are in equilibrium if a change in strategies by any one of them would lead that player to earn less than if he remained with his current strategy. A *Nash equilibrium* satisfies the following condition

$$E(p, p) \geq E(r, p). \quad (1)$$

A player can not increase his payoff if he decides to play the strategy r instead of p .

Each agent in a n -player game where the i^{th} player has as strategy space S_i is modeled by a population of players which have to be partitioned into groups. Individuals in the same group would all play the same strategy. Randomly we make play the members of the subpopulations against each other. The subpopulations that perform the best will grow and those that do not will shrink and eventually will vanish. The process of natural selection assures survival of the best players at the expense of the others. A population equilibrium occurs when the population shares are such that the expected payoffs for all strategies are equal.

Consider a large population in which a two person game $G = (S, E)$ is played by randomly matched pairs of animals generation after generation. Let p be the strategy played by the vast majority of the population, and let r be the strategy of a mutant present in small frequency. Both p and r can be pure or mixed. An *evolutionary stable strategy* (ESS) p of a symmetric two - person game $G = (S, E)$ is a pure or mixed strategy for G which satisfies the following two conditions

$$\begin{aligned} E(p, p) &> E(r, p), \\ \text{If } E(p, p) &= E(r, p) \text{ then } E(p, r) > E(r, r). \end{aligned} \quad (2)$$

Since the stability condition only concerns to alternative best replies, p is always evolutionarily stable if (p, p) is an strict equilibrium point. An ESS is also a Nash equilibrium since it is the best reply to itself and the game is symmetric. The set of all the strategies that are ESS is a subset of the NE of the game. A population which plays an ESS can withstand an invasion by a small group of mutants playing a different strategy. It means that if

a few individuals which play a different strategy are introduced into a population in an ESS, the evolutionarily selection process would eventually eliminate the invaders.

The natural selection process that determines how populations playing specific strategies evolve is known as the *replicator dynamics*. It describes the evolution of a polymorphic state in a population represented by a mixed strategy x for G whose members are involved in a conflict described by a symmetric two - person game $G = (S, E)$. The probability assigned to a pure strategy s is denoted by $x(s)$. If $s_i, i = 1, \dots, n \in S$ are the pure strategies available to a player, then that player's strategy will be denoted by the column vector x where $x_i \in [0, 1]$ and $\sum_{i=1}^n x_i = 1$ in which the i^{th} component of x gives the probability of playing strategy s_i . The i^{th} component of x also is interpreted as the relative frequency of individuals using strategy s_i . Playing a pure strategy s_j is represented by the vector x whose j^{th} component is 1, and all the other components are 0. The fitness function is given by $E = f_i(x)$, $i = 1, \dots, n$ and specifies how successful each subpopulation is. The fitness function must be defined for each component of x . The fitness for x_i is the expected utility of playing strategy s_i against a player with a mixed strategy defined by the vector x and is given by

$$f_i(x) = (Ax)_i, \quad (3)$$

where A is the payoff matrix and the subscript $i = 1, \dots, n$ denotes the i^{th} component of the matrix-vector product. Or also

$$f_i(x) = \sum_{j=1}^n a_{ij}x_j, \quad (4)$$

where a_{ij} are the elements of the payoff matrix A . The average fitness of the population is given by

$$\langle f(x) \rangle = x^T Ax, \quad (5)$$

$$\langle f(x) \rangle = \sum_{i=1}^n x_i f_i(x), \quad (6)$$

$$\langle f(x) \rangle = \sum_{k,l=1}^n a_{kl}x_kx_l. \quad (7)$$

The superscript T denotes transpose.

The evolution of relative frequencies in a population is described by the replicator dynamics

$$\frac{dx_i(t)}{dt} = [f_i(x) - \langle f(x) \rangle] x_i(t). \quad (8)$$

By replacing relations (3) and (5) in equation (8) and by replacing relations (4) and (7) we can obtain two different

ways of representing the replicator dynamics

$$\frac{dx_i(t)}{dt} = [(Ax)_i - x^T Ax] x_i(t), \quad (9)$$

$$\frac{dx_i(t)}{dt} = \left[\sum_{j=1}^n a_{ij}x_j - \sum_{k,l=1}^n a_{kl}x_kx_l \right] x_i(t). \quad (10)$$

The stable fixed points of the replicator dynamics are Nash equilibria. It is important to note that the fixed points of a system do not change in the time. It means that if a population reaches a state which is a Nash equilibrium, it will remain there. There are different versions of these equations [5, 6, 9, 11] depending on the evolutionary model used. The replicator dynamics rewards strategies that outperform the average by increasing their frequency, and penalizes poorly performing strategies by decreasing their frequency.

In a symmetric game payoff matrices and actions are identical for both agents. These games can be modeled by a single population of individuals playing against each other. When the game being played is asymmetric, a different population of players must be used to simulate each agent. For a two-player asymmetric normal game each player will have a distinct payoff matrix A and B . The strategy vector for player one is represented by x and for player two is represented by y . Player one has n strategies $s_{1i}, i = 1, \dots, n \in S_1$ and player two has m strategies $s_{2j}, j = 1, \dots, m \in S_2$. The fitness for a player who plays the strategy s_{1i} will be $f_{1i} = (Ay)_i$ and the average fitness of the first population will be $\langle f_1 \rangle = x^T Ay$. The fitness for a player who plays the strategy s_{2i} will be $f_{2i} = (Bx)_i$ and the average fitness of the second population will be $\langle f_2 \rangle = y^T Bx$. The evolution of this game would be described for the next equations system

$$\begin{aligned} \frac{dx_i(t)}{dt} &= [(Ay)_i - x^T Ay] x_i(t), \\ \frac{dy_i(t)}{dt} &= [(Bx)_i - y^T Bx] y_i(t). \end{aligned} \quad (11)$$

The genetic code is the relationship between the sequence of the bases in the DNA and the sequence of amino acids in proteins. Recent work [42] about evolvability of the genetic code suggests that the code is shaped by natural selection.

III. REPLICATOR DYNAMICS MATRIX FORM

As we saw the replicator dynamics is a differential equation where x is a column vector. Obviously, the matrix $U = (Ax)_i - x^T Ax$ has to be diagonal and its elements are given by

$$u_{ii} = \sum_{j=1}^n a_{ij}x_j - \sum_{k,l=1}^n a_{kl}x_kx_l. \quad (12)$$

The replicator dynamics can be expressed as

$$\frac{dx}{dt} = Ux. \quad (13)$$

By multiplying each element of the vector x by its corresponding $(x_i)^{-1/2}$ in both parts of equation (10) or (13), we can get

$$v = U\hat{x}, \quad (14)$$

where v and $\hat{x}$ are column vectors with elements $v_i = \frac{1}{(x_i)^{1/2}} \frac{dx_i}{dt}$ and $\hat{x}_i = (x_i)^{1/2}$ respectively. Lets multiply equation (14) by $\hat{x}^T$ and lets define the matrix

$$G = \frac{1}{2}v\hat{x}^T, \quad (15)$$

where $g_{ij} = \frac{1}{2} \frac{(x_j)^{1/2}}{(x_i)^{1/2}} \frac{dx_i}{dt}$ are the elements of G . The matrix G is also equal to

$$G = \frac{1}{2}U\hat{x}\hat{x}^T \quad (16)$$

with

$$(U\hat{x}\hat{x}^T)_{ij} = \left(\sum_{k=1}^n a_{ik}x_k - \sum_{k,l=1}^n a_{kl}x_kx_l \right) (x_ix_j)^{1/2}.$$

Lets find G^T

$$\begin{aligned} G^T &= \frac{1}{2}(v\hat{x}^T)^T, \\ G^T &= \frac{1}{2}\hat{x}v^T, \end{aligned} \quad (17)$$

where $g_{ij}^T = \frac{1}{2} \frac{(x_i)^{1/2}}{(x_j)^{1/2}} \frac{dx_j}{dt}$ are the elements of G^T . The matrix G^T is also equal to

$$\begin{aligned} G^T &= \frac{1}{2}(U\hat{x}\hat{x}^T)^T, \\ G^T &= \frac{1}{2}\hat{x}\hat{x}^T U^T \end{aligned} \quad (18)$$

with

$$(\hat{x}\hat{x}^T U^T)_{ij} = (x_jx_i)^{1/2} \left(\sum_{k=1}^n a_{jk}x_k - \sum_{k,l=1}^n a_{kl}x_kx_l \right).$$

Lets define

$$\frac{dX}{dt} = G + G^T \quad (19)$$

so that the matrix X has as elements

$$x_{ij} = (x_ix_j)^{1/2}. \quad (20)$$

The time evolution of X is also equal to

$$G + G^T = \frac{1}{2}(U\hat{x}\hat{x}^T + \hat{x}\hat{x}^T U^T) \quad (21)$$

with

$$\begin{aligned} (G + G^T)_{ij} &= \frac{1}{2} \sum_{k=1}^n a_{ik}x_k (x_ix_j)^{1/2} \\ &\quad + \frac{1}{2} \sum_{k=1}^n a_{jk}x_k (x_jx_i)^{1/2} \\ &\quad - \sum_{k,l=1}^n a_{kl}x_kx_l (x_ix_j)^{1/2}. \end{aligned} \quad (22)$$

Lets call

$$(G_1)_{ij} = \frac{1}{2} \sum_{k=1}^n a_{ik}x_k (x_ix_j)^{1/2}, \quad (23)$$

$$(G_2)_{ij} = \frac{1}{2} \sum_{k=1}^n a_{jk}x_k (x_jx_i)^{1/2}, \quad (24)$$

$$(G_3)_{ij} = \sum_{k,l=1}^n a_{kl}x_kx_l (x_ix_j)^{1/2} \quad (25)$$

the elements of the matrixes G_1 , G_2 and G_3 that compose by adding the matrix $(G + G^T)$. We can factorize G_1 in two matrixes Q and X . It means

$$G_1 = QX, \quad (26)$$

where Q is a diagonal matrix and has as elements $q_{ii} = \frac{1}{2} \sum_{k=1}^n a_{ik}x_k$. In the same way

$$G_2 = XQ. \quad (27)$$

The matrix G_3 can also be represented as $(G_3)_{ij} = \sum_{l=1}^n (x_ix_l)^{1/2} \sum_{k=1}^n a_{kl}x_k (x_lx_j)^{1/2}$ and it can be factorized into

$$G_3 = 2XQX. \quad (28)$$

It is easy to show that $X^2 = X$ so that we can write the equation (19) like

$$\frac{dX}{dt} = QXX + XXQ - 2XQX \quad (29)$$

and finally, by grouping into commutators

$$\frac{dX}{dt} = [[Q, X], X], \quad (30)$$

$$\frac{dX(t)}{dt} = [\Lambda(t), X(t)], \quad (31)$$

where $\Lambda = [Q, X]$.

The elements of Λ are given by $(\Lambda)_{ij} = \frac{1}{2} [(\sum_{k=1}^n a_{ik}x_k)(x_ix_j)^{1/2} - (x_jx_i)^{1/2}(\sum_{k=1}^n a_{jk}x_k)]$. Equations (30) and (31) can be considered as a generalization of the replicator dynamics as was seen in equations (8), (9) and (10). We could call X the relative frequency matrix and its evolution is described by equation (31). Matrix X has the following properties

- a) $Tr(X) = 1$,
- b) $X^2 = X$,
- c) $X^T = X$.

It is easy to realize that each component of this matrix will evolve following the replicator dynamics as was seen

in equations (8), (9) and (10) so that we could call equation (31) the matrix form of the replicator dynamics. If we take $\Theta = [\Lambda, X]$ equation (31) becomes into

$$\frac{dX}{dt} = \Theta, \quad (32)$$

where the elements of Θ are given by

$$\begin{aligned} (\Theta)_{ij} = & \frac{1}{2} \sum_{k=1}^n a_{ik} x_k (x_i x_j)^{1/2} \\ & + \frac{1}{2} \sum_{k=1}^n a_{jk} x_k (x_j x_i)^{1/2} \\ & - \sum_{k,l=1}^n a_{lk} x_k x_l (x_i x_j)^{1/2}. \end{aligned} \quad (33)$$

The fact that we can calculate the evolution of interference effects between relative frequencies like $(x_i x_j)^{1/2}$ is the advantage of using equation (31) besides equation (10).

IV. MIXED AND PURE STATES IN QUANTUM MECHANICS

In classical mechanics we can precisely specify the state of a system by one point in its phase space. Its trajectory through the phase space describes the time evolution of the system and this evolution follows Newton's laws or Hamilton equations. In quantum mechanics we can precisely describe the state of a system by specifying its state vector $|\Psi(t)\rangle$ or its wave function $\Psi(r, t)$ and the evolution of the system is given by Schrödinger equation. However, in most cases when the information about the system is incomplete the state of a system is not perfectly defined for which we have to describe our system in terms of probabilities [43, 44].

An *ensemble* is a collection of identically prepared physical systems. When each member of the ensemble is characterized by the same state vector $|\Psi(t)\rangle$ it is called *pure ensemble*. If each member has a probability p_i of being in the state $|\Psi_i(t)\rangle$ we have a *mixed ensemble*. The state of a system $|\Psi_i(t)\rangle = \left| \Psi_k^{(i)}(t) \right\rangle$ is a column vector $\left| \Psi_k^{(i)}(t) \right\rangle = (\Psi_1^{(i)}(t), \Psi_2^{(i)}(t), \dots, \Psi_n^{(i)}(t))^T, k = 1, \dots, n$.

For the case of *pure ensemble* the expected value of measure certain observable described by the operator A in the state $|\Psi(t)\rangle$ at the instant t is given by

$$\begin{aligned} \langle A \rangle &= \langle \Psi(t) | A | \Psi(t) \rangle = \sum_{i,j=1}^n \langle j | \Psi(t) \rangle a_{ij} \langle \Psi(t) | i \rangle, \\ \langle A \rangle &= \sum_{i,j=1}^n a_{ij} c_i^*(t) c_j(t), \end{aligned} \quad (34)$$

where a_{ij} are the elements of the matrix that represents the observable A . The terms $c_i^*(t) = \langle \Psi(t) | i \rangle$ and $c_j(t) = \langle j | \Psi(t) \rangle$ are the elements of certain operator $\rho(t)$

$$c_i^*(t) c_j(t) = \langle j | \Psi(t) \rangle \langle \Psi(t) | i \rangle = \langle j | \rho(t) | i \rangle \quad (35)$$

defined as

$$\rho(t) = |\Psi(t)\rangle \langle \Psi(t)|. \quad (36)$$

The density operator $\rho(t)$ for a pure ensemble in a state $|\Psi(t)\rangle$ satisfies the following properties

- a) $\rho^+(t) = \rho(t)$,
- b) $\rho^2(t) = \rho(t)$,
- c) $Tr \rho^2(t) = 1$.

The superscript $+$ denotes Hermitian. We can make all the predictions made for $|\Psi(t)\rangle$ in function of $\rho(t)$. The sum of the diagonal elements of the density matrix is one

$$\sum_n |c_n(t)|^2 = Tr \rho(t) = 1. \quad (37)$$

The mean value of an observable A is calculated using

$$\langle A \rangle = Tr \{ \rho(t) A \}. \quad (38)$$

The time evolution of the density operator is given by the following equation

$$i\hbar \frac{d\rho(t)}{dt} = [H(t), \rho(t)]. \quad (39)$$

Each member of a *mixed ensemble* is a pure state and its evolution is given by Schrödinger equation. The probabilities for each state are constrained to satisfy the normalization condition

$$\sum_{i=1}^n p_i = 1, \quad (40)$$

where $0 \leq p_1, p_2, \dots, p_n \leq 1$. Suppose we make a measurement on a mixed ensemble of some observable A . The *ensemble average* of A is defined by the average of the expected values measured in each member of the ensemble described by $|\Psi_i(t)\rangle$ and with probability p_i

$$\langle A \rangle_\rho = \sum_{i=1}^n p_i \langle \Psi_i(t) | A | \Psi_i(t) \rangle \quad (41)$$

it means $\langle A \rangle_\rho = p_1 \langle A \rangle_1 + p_2 \langle A \rangle_2 + \dots + p_n \langle A \rangle_n$.

$$\begin{aligned} \langle A \rangle_\rho &= \sum_{i=1}^n p_i \langle \Psi_i(t) | A | \Psi_i(t) \rangle, \\ \langle A \rangle_\rho &= \sum_{i,j,k=1}^n p_i \langle \Psi_i(t) | j \rangle \langle j | A | k \rangle \langle k | \Psi_i(t) \rangle, \\ \langle A \rangle_\rho &= \sum_{i,j,k=1}^n p_i a_{jk} c_j^{(i)*}(t) c_k^{(i)}(t), \end{aligned} \quad (42)$$

where a_{jk} are the elements of the matrix that represents the observable A . The terms $c_k^{(i)}(t) = \langle k | \Psi_i(t) \rangle$ and $c_j^{(i)*}(t) = \langle \Psi_i(t) | j \rangle$ are the elements of certain operator $\rho(t)$

$$\sum_{i=1}^n p_i c_j^{(i)*}(t) c_k^{(i)}(t) = \sum_{i=1}^n \langle k | \Psi_i(t) \rangle p_i \langle \Psi_i(t) | j \rangle, \quad (43)$$

$$\sum_{i=1}^n p_i c_j^{(i)*}(t) c_k^{(i)}(t) = \langle k | \rho(t) | j \rangle$$

now defined as

$$\rho(t) = \sum_{i=1}^n p_i |\Psi_i(t)\rangle \langle \Psi_i(t)|. \quad (44)$$

The density operator contains all the physically significant information we can possibly obtain about the ensemble in question. Any two ensembles that produce the same density operator are physically indistinguishable. From equation (42) we have

$$\langle A \rangle = \text{Tr} \{ \rho(t) A \}. \quad (45)$$

The density operator for a mixed ensemble satisfies the next properties

- a) ρ is Hermitian,
- b) $\text{Tr} \rho(t) = 1$,
- c) $\rho^2(t) \leq \rho(t)$,
- d) $\text{Tr} \rho^2(t) \leq 1$.

A pure state is specified by $p_i = 1$ for some $|\Psi_i(t)\rangle$, $i = 1, \dots, n$ and the density operator $\rho(t)$ is represented by a matrix with all its elements equal to zero except one 1 on the diagonal.

The diagonal elements ρ_{nn} of the density operator $\rho(t)$ represents the average probability of finding the system in the state $|n\rangle$

$$\rho_{nn} = \langle n | \rho(t) | n \rangle = \sum_{i=1}^n \langle n | \Psi_i(t) \rangle p_i \langle \Psi_i(t) | n \rangle, \quad (46)$$

$$\rho_{nn} = \sum_{i=1}^n p_i \left| c_n^{(i)} \right|^2,$$

where $c_n^{(i)} = \langle n | \Psi_i(t) \rangle$ and $\left| c_n^{(i)} \right|^2 \in \mathbb{R}^+$. If the state of the system is $|\Psi_i(t)\rangle$, $\left| c_n^{(i)} \right|^2$ is the probability of finding, in a measurement, this system in the state $|n\rangle$. The diagonal elements ρ_{nn} are zero if and only if all $\left| c_n^{(i)} \right|^2$ are zero. The non-diagonal elements ρ_{np} expresses the interference effects between the states $|n\rangle$ and $|p\rangle$ which can

appear when the state $|\Psi_i\rangle$ is a coherent linear superposition of these states

$$\rho_{np} = \langle n | \rho(t) | p \rangle = \sum_{i=1}^n \langle n | \Psi_i(t) \rangle p_i \langle \Psi_i(t) | p \rangle, \quad (47)$$

$$\rho_{np} = \sum_{i=1}^n p_i c_n^{(i)}(t) c_p^{(i)*}(t)$$

with $c_n^{(i)}(t) = \langle n | \Psi_i(t) \rangle$, $c_p^{(i)*}(t) = \langle \Psi_i(t) | p \rangle$ and $c_n^{(i)} c_p^{(i)*} \in \mathbb{C}$. If $\rho_{np} = 0$, it means that the average has canceled out any interference effects between $|n\rangle$ and $|p\rangle$ but if it is different from zero subsists certain coherence between these states.

The time evolution of the density operator that describes a mixed or pure ensemble is given by

$$i\hbar \frac{d\rho(t)}{dt} = [H(t), \rho(t)] \quad (48)$$

which is the quantum analogue of Liouville's theorem from classical mechanics and a generalization of Schrödinger equation.

V. DENSITY OPERATOR & QUANTUM STATISTICAL MECHANICS

Now lets review briefly the connection between the density operator and quantum statistical mechanics [44]. The basic assumption we make to obtain density operator from the study of an ensemble in thermal equilibrium is that nature tends to maximize σ subject to the constraint that the ensemble average of the Hamiltonian has a certain prescribed value. The quantity σ is defined by

$$\sigma = -\text{Tr}(\rho \ln \rho) \quad (49)$$

which can be regarded as a quantitative measure of disorder.

It is easy to show that for a completely random ensemble in where the density matrix diagonalized looks like

$$\rho = \frac{1}{N} \begin{pmatrix} 1 & & & & 0 \\ & 1 & & & \\ & & 1 & & \\ & & & \dots & \\ & & & & 1 \\ 0 & & & & & 1 \end{pmatrix} \quad (50)$$

the quantity σ takes the value

$$\sigma = \ln N \quad (51)$$

and in contrast, for a pure ensemble with a density matrix diagonalized similar to

$$\rho = \begin{pmatrix} 0 & & & & 0 \\ & 0 & & & \\ & & 1 & & \\ & & & 0 & \\ & & & \dots & 0 \\ 0 & & & & 0 \end{pmatrix} \quad (52)$$

the quantity σ takes the value

$$\sigma = 0. \quad (53)$$

A pure ensemble is an ensemble with a maximum amount of order because all members are characterized by the same quantum mechanical state ket. By the other hand, a completely random ensemble in which all quantum mechanical states are equally likely σ takes its maximum possible value subject to the normalization condition

$$\sum_k \rho_{kk} = 1. \quad (54)$$

Our quantity σ is related to the definition of entropy S from quantum statistical mechanics via

$$S = k\sigma, \quad (55)$$

where k is the Boltzmann constant. Once that σ is a maximum and thermal equilibrium is established, we expect

$$\frac{d\rho}{dt} = 0. \quad (56)$$

We maximize σ by requiring that

$$\delta\sigma = 0 \quad (57)$$

subject to the constrains

$$\delta \langle H \rangle = \sum_k \delta\rho_{kk} E_k = 0 \quad (58)$$

and

$$\delta(Tr(\rho)) = \sum_k \delta\rho_{kk} = 0. \quad (59)$$

By the use of Lagrange multipliers we can accomplish these and we can obtain

$$\sum_k \delta\rho_{kk} [(\ln \rho_{kk} + 1) + \beta E_k + \gamma] = 0 \quad (60)$$

which for an arbitrary variation is possibly only if

$$\rho_{kk} = \exp(-\beta E_k - \gamma - 1), \quad (61)$$

where we can eliminate γ using the normalization condition and our final result is

$$\rho_{kk} = \frac{\exp(-\beta E_k)}{\sum_l \exp(-\beta E_l)}. \quad (62)$$

The denominator of equation (62) is the partition function

$$Z = \sum_l \exp(-\beta E_l) = Tr(e^{-\beta H}). \quad (63)$$

Given ρ_{kk} in the energy basis we can write the density operator as

$$\rho = \frac{e^{-\beta H}}{Z}. \quad (64)$$

The operator ρ is the quantum mechanical analogue of the equilibrium density of points in the phase space for an canonical ensemble.

The diagonal elements of the density operator in the $\{|n\rangle\}$ basis of eigenvectors of H are given by

$$\rho_{nn} = \frac{\langle n | e^{-\beta H} | n \rangle}{Z} = \frac{e^{-\beta E_n}}{Z} \quad (65)$$

and the non diagonal elements are equal to

$$\rho_{np} = \frac{\langle n | e^{-\beta H} | p \rangle}{Z} = \frac{e^{-\beta E_p}}{Z} \langle n | p \rangle = 0. \quad (66)$$

At thermodynamic equilibrium, the populations of the stationary states are exponentially decreasing functions of the energy, and the coherences between stationary states are zero [43].

VI. QUANTIZATION RELATIONSHIPS & QUANTUM REPLICATOR DYNAMICS

Let us propose the next quantization relationships

$$x_i \rightarrow \sum_{k=1}^n \langle i | \Psi_k \rangle p_k \langle \Psi_k | i \rangle = \rho_{ii}, \quad (67)$$

$$(x_i x_j)^{1/2} \rightarrow \sum_{k=1}^n \langle i | \Psi_k \rangle p_k \langle \Psi_k | j \rangle = \rho_{ij}. \quad (68)$$

It means that a population will be represented by a quantum system in which each subpopulation playing strategy s_i will be represented by a pure ensemble in the state $|\Psi_k(t)\rangle$ and with probability p_k . The probability x_i of playing strategy s_i or the relative frequency of the individuals using strategy s_i in that population will be represented as the probability ρ_{ii} of finding each pure ensemble in the state $|i\rangle$. Our matrix X of relative frequencies of a population corresponds to the density matrix ρ

$$X \rightarrow \rho \quad (69)$$

and from equations (31) and (48)

$$\Lambda \rightarrow \hat{H}, \quad (70)$$

where $H = i\hbar\hat{H}$. Through these quantization relationships the replicator dynamics takes the form of the equation of evolution of mixed states

$$i\hbar\frac{d\rho(t)}{dt} = [H(t), \rho(t)] \quad (71)$$

and according with equation (49) the entropy of our system would be given by

$$\sigma = -Tr(X \ln X), \quad (72)$$

where x_{ij} are the elements of the matrix X .

We can translate our classical definitions to the quantum world like the payoff function

$$f_i(\rho) = \sum_{j=1}^n a_{ij}\rho_{jj} \quad (73)$$

and the average fitness of the population

$$\langle f(\rho) \rangle = \sum_{k,l=1}^n a_{kl}\rho_{kk}\rho_{ll}. \quad (74)$$

When the non diagonal elements of A and ρ are zero

$$Q = \frac{1}{2}A\rho \quad (75)$$

so that we can rewrite Λ as

$$\Lambda = \frac{1}{2}[A\rho - \rho A\rho] \quad (76)$$

and due to the fact that $H = i\hbar\Lambda$

$$H = \frac{i\hbar}{2}[A\rho - \rho A\rho] \quad (77)$$

the average energy of our system is

$$\begin{aligned} \langle E \rangle &= Tr(\rho H), \\ \langle E \rangle &= \frac{i\hbar}{2}Tr[\rho A\rho - \rho A\rho] = 0. \end{aligned} \quad (78)$$

For this case it is easy to note in equation (76) that the Hamiltonian H of our system is a diagonal matrix in where its elements H_{ii} correspond to its eigenvalues and its eigenvectors are

$$\begin{aligned} |\Psi_1\rangle &= c_{11}|1\rangle + c_{12}|0\rangle + \dots + c_{1i}|0\rangle + \dots + c_{1n}|0\rangle, \\ |\Psi_2\rangle &= c_{21}|0\rangle + c_{22}|1\rangle + \dots + c_{2i}|0\rangle + \dots + c_{2n}|0\rangle, \\ &\dots \\ |\Psi_n\rangle &= c_{n1}|0\rangle + c_{n2}|0\rangle + \dots + c_{ni}|0\rangle + \dots + c_{nn}|1\rangle \end{aligned} \quad (79)$$

subject to the constrain $\sum_{j=1}^n |c_{ij}|^2 = 1$. For the case of a two person symmetric game with $a_{ij} = 0$ and $\rho_{ij} = 0$ the eigenvectors of the game are

$$\begin{aligned} |\Psi_1\rangle &= \alpha|1\rangle + \beta|0\rangle, \\ |\Psi_2\rangle &= \gamma|0\rangle + \delta|1\rangle, \end{aligned} \quad (80)$$

where $\alpha, \beta, \gamma, \delta$ are subject to the constraints $|\alpha|^2 + |\beta|^2 = 1$ and $|\gamma|^2 + |\delta|^2 = 1$. For this case also $[\Lambda, \rho] = \Lambda$ and

$$\frac{dX}{dt} = \Lambda. \quad (81)$$

The payoff function takes the form $f(x) = A\rho$ and its trace is equal to

$$Tr(f(x)) = \langle A \rangle_\rho \quad (82)$$

and the average payoff function is given by

$$\langle f(x) \rangle = Tr(\rho A\rho). \quad (83)$$

For $x_{ij} = 0$ our entropy takes the form

$$\sigma = -\sum_{i=1}^n x_{ii} \ln x_{ii} = -\sum_{i=1}^n x_i \ln x_i. \quad (84)$$

Finally, the equation of evolution of mixed states from quantum statistical mechanics (48) is the quantum analogue of the replicator dynamics in matrix form (31) and the respective matrixes of both systems have similar properties.

VII. QUANTUM STRATEGIC EQUILIBRIUM

Quantum mechanics could be a much more general theory that we had thought. It could encloses theories like EGT and evolutionary dynamics and we could explain through this theory biological and economical processes. From this point of view many of the equations, concepts and its properties defined quantically must be more general than its classical versions which must remain inside of the foundations of the new quantum version. So, our quantum equilibrium concept also must be more general than the one defined classically.

If in an isolated system each of its accessible states do not have the same probability, the system is not in equilibrium. The system will vary and will evolution in the time until it reaches the equilibrium state when the probability of finding the system in each of the accessible states be the same. Then the system will find its more probable configuration in which the number of accessible states is maximum and equally probable. The whole system will vary and rearrange its state and the states of its ensembles with the purpose of maximize its entropy and reach its minimum energy state. We could say that the purpose and maximum payoff of a quantum system is its minimum energy state.

The system and its members will vary and rearrange themselves to reach the best possible state for each of them which is also the best possible state for the whole system. This can be seen like a microscopical cooperation between quantum objects to improve its state with the

purpose of reaching or maintaining the equilibrium of the system. All the members of our quantum system will play a game in which its maximum payoff is its minimum energy state. Different particles will cooperate to form a system in which its new state will have to be better than the one they had before they interact by forming an atom. Different atoms will cooperate to form a system in which its new state have to be better than the one they had before they formed what is called a “inorganic” molecule.

A system is stable only if it maximizes the welfare of the collective above the welfare of the individual, if it is maximized the welfare of the individual above the welfare of the collective the system gets unstable and eventually it collapses.

Gafiychuk and Prykarpatsky [45] applied the replicator equations written in the form of nonlinear von Neumann equations to the study of the general properties of the quasispecies dynamical system from the standpoint of its evolution and stability. They developed a mathematical model of a naturally fitted coevolving ecosystem and a theoretical study a self-organization problem of an ensemble of interacting species. DNA is a nonlinear dynamical system and its evolution is a sequence of chemical reactions. An abstract DNA-type system is defined by a set of nonlinear kinetic equations with polynomial nonlinearities that admit soliton solutions associated with helical geometry. Aerts and Czachor [46] shown that the set of these equations allows for two different Lax representations: They can be written as von Neumann type nonlinear systems and they can be regarded as a compatibility condition for a Darboux-covariant Lax pair. Organisms whose DNA evolves in a chaotic way would be eliminated by natural selection. They also explained why non-Kolmogorovian probability models occurring in soliton kinetics are naturally associated with chemical reactions. Gogonea and Merz [38] indicated that games are being played at the quantum mechanical level in protein folding. Turner and Chao [47] studied the evolution of competitive interactions among viruses in an RNA phage, and found that the fitness of the phage generates a payoff matrix conforming to the two-person prisoner’s dilemma game. Bacterial infections by viruses have been presented as classical game-like situations where nature prefers the dominant strategies. Patel [39, 40] suggested

quantum dynamics played a role in the DNA replication and the optimization criteria involved in genetic information processing. He considers the criteria involved as a task similar to an unsorted assembly operation where the Grover’s database search algorithm fruitfully applies; given the different optimal solutions for classical and quantum dynamics. Azhar Iqbal [22] showed results in which quantum mechanics has strong and important roles in selection of stable solutions in a system of interacting “entities”. These entities can do quantum actions on quantum states. It may simply consists of a collection of molecules and the stability of solutions or equilibria can be affected by quantum interactions which provides a new approach towards theories of rise of complexity in groups of quantum interacting entities. Neuroeconomics [48, 49] may provide an alternative to the classical Cartesian model of the brain and behavior [50] through a rich dialogue between theoretical neurobiology and quantum logic [51, 52].

VIII. CONCLUSIONS

It has been shown that through the quantization relationships proposed the quantum analogue of the replicator dynamics is the equation of evolution of mixed states of a quantum system. A population is represented by a quantum system in which each subpopulation playing strategy s_i is represented by a pure ensemble in state $|\Psi_k\rangle$ and with probability p_k . The probability x_i of playing strategy s_i or the relative frequency of the individuals using strategy s_i in that population will be represented as the probability ρ_{ii} of finding each pure ensemble in the state $|i\rangle$. Through quantum mechanics we can describe economical and biological processes from a different perspective and with a quantum equilibrium concept which is more general than whichever defined classically. The purpose and maximum payoff of a system is improve its state by maximizing the welfare of its members through maximizing the welfare of the system. For a quantum system the maximum payoff is its minimum energy state. The system and all its members will cooperate and rearrange its states to improve its present condition. A system is stable only if it maximizes the welfare of the collective above the welfare of the individual or else the system gets unstable and eventually it collapses.

-
- [1] J. von Neumann and O. Morgenstern, *The Theory of Games and Economic Behavior* (Princeton University Press, Princeton, 1947).
 - [2] R. B. Myerson, *Game Theory: An Analysis of Conflict* (MIT Press, Cambridge, 1991).
 - [3] M. A. Nowak and K. Sigmund, *Nature* **398**, 367 (1999).
 - [4] J. M. Smith, *Evolution and The Theory of Games* (Cambridge University Press, Cambridge, UK, 1982).
 - [5] J. Hofbauer and K. Sigmund, *Evolutionary Games and Replicator Dynamics* (Cambridge University Press, Cambridge, UK, 1998).
 - [6] J. Weibull, *Evolutionary Game Theory* (MIT Press, Cambridge, MA, 1995).
 - [7] R. A. Fisher, *The Genetic Theory of Natural Selection* (Oxford, Clarendon Press, 1930).
 - [8] P. Hammerstein and R. Selten, *Game Theory and Evo-*

- lutionary Biology* (Handbook of Game Theory. Vol 2. Elsevier B.V., 1994).
- [9] R. Cressman, *The Stability Concept of Evolutionary Game Theory: A Dynamic Approach* (Springer-Verlag, New York, 1992).
 - [10] J. M. Smith and G. R. Price, *The logic of animal conflict*, Nature **246**, 15 (1973).
 - [11] P. D. Taylor and L. B. Jonker, *Evolutionary stable strategies and game dynamics*, Mathematical Biosciences **40**, 145-156 (1978).
 - [12] D. A. Meyer, Phys. Rev. Lett. **82**, 1052-1055 (1999).
 - [13] J. Eisert, M. Wilkens and M. Lewenstein, Phys. Rev. Lett. **83**, 3077 (1999).
 - [14] J. Eisert and M. Wilkens, J. Mod. Opt. **47**, 2543 (2000).
 - [15] L. Marinatto and T. Weber, Phys. Lett. A **272**, 291 (2000).
 - [16] C. F. Lee and N. F. Johnson, Phys. Rev. A **67**, 022311 (2003).
 - [17] J. Du, H. Li, X. Xu, M. Shi, J. Wu, X. Zhou, and R. Han, Phys. Rev. Lett. **88**, 137902 (2002).
 - [18] A. P. Flitney and D. Abbott, Proc. R. Soc. (London) A **459**, 2463-74 (2003).
 - [19] J. Shimamura, S. K. Ozdemir, F. Morikoshi and N. Imoto, Int. J. of Quant. Inf. **2/1**, 79 (2004).
 - [20] S. K. Ozdemir, J. Shimamura, F. Morikoshi and N. Imoto, Phys. Lett. A **333**, 218 (2004).
 - [21] E. W. Piotrowski and J. Sladkowski, Int. J. Theor. Phys. **42**, 1089 (2003).
 - [22] Azhar Iqbal, PhD thesis, Quaid-i-Azam University, 2004, quant-ph/0503176.
 - [23] S. C. Benjamin and P. M. Hayden, Phys. Rev. A **64**, 030301 (2001).
 - [24] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
 - [25] C. F. Lee and N. F. Johnson, *Parrondo Games and Quantum Algorithms*, quant-ph/0204043.
 - [26] P. Moulin and A. Ivanovic, Proc. of Int. Conf. on Image Process. **3**, 975-978 (2001).
 - [27] A. S. Cohen and A. Lapidot, IEEE Trans. on Inf. Theory **48**, 1639-1667 (2002).
 - [28] J. Conway and N. Sloane, IEEE Trans. on Inf. Theory **32**, 337-348 (1986).
 - [29] X. M. Shen and L. Deng, IEEE Trans. on Signal Process. **45**, 1092-1095 (1997).
 - [30] J. M. Ettinger, *2nd Inf. Hiding Workshop*, Portland, OR, Apr 15-17 (1998).
 - [31] S. Pateux and G. Le Guelvouit, Elsevier, Signal Process. Image Comm. **18**, 283-296 (2003).
 - [32] L. A. DaSilva, and V. Srivastava, *The First Workshop on Games and Emergent Behaviors in Distributed Computing Environments*, Birmingham, UK, September (2004).
 - [33] E. W. Piotrowski and J. Sladkowski, *Quantum Market Games*, Physica A **312**, 208-216, (2002).
 - [34] E. W. Piotrowski and J. Sladkowski, *Quantum-like Approach to Financial Risk: Quantum Anthropic Principle*, Acta Phys. Pol. B **32**, 3873-3879, (2001).
 - [35] E. W. Piotrowski and J. Sladkowski, *Interference of quantum market strategies*, Physica A **318**, 516-518 (2003).
 - [36] R. Dawkins, *The Selfish Gene* (Oxford University Press, Oxford, 1976).
 - [37] R. Axelrod, *The Evolution of Cooperation* (Basic Books, New York, 1984).
 - [38] V. Gogonea and K. M. Merz, *Fully quantum mechanical description of proteins in solution – combining linear scaling quantum mechanical methodologies with the Poisson-Boltzmann equation*, J. Phys. Chem. A **103**, 5171-5188 (1999).
 - [39] A. Patel, *Quantum algorithms and the genetic code*, Pragma **56**, 367-381, (2001).
 - [40] A. Patel, *Testing Quantum Dynamics in Genetic Information Processing*, Journal of Genetics **80**, 39, (2001).
 - [41] D. Home and R. Chattopadhyaya, *Determination of When an Outcome is Actualised in a Quantum Measurement using DNA — Photolyase System*, quant-ph/9903036.
 - [42] R. D. Knight, S.J. Freeland, and L.F. Landweber, *Rewiring the keyboard: evolvability of genetic code*, Nature Reviews Genetics **2**, 49, (2001).
 - [43] C. Tanoudji and B. Diu, Lao, *Quantum Mechanics* (Herman, Paris, 1977).
 - [44] J. J. Sakurai, *Modern Quantum Mechanics*, (Addison - Wesley, 1994).
 - [45] V. Gafychuk and A. Prykarpatsky, J. Nonlin. Math Phys. **11**, 350 (2004).
 - [46] D. Aerts and M. Czachor, *Abstract DNA-type systems*, q-bio.PE/0411031.
 - [47] P. E. Turner and L. Chao, Prisoner's dilemma in an RNA virus, Nature **398**, 6726 (1999).
 - [48] P. W. Glimcher, *Decisions, Decisions, Decisions: Choosing a Biological Science of Choice*, Neuron **36**, 323 (2002).
 - [49] P. W. Glimcher, *Decisions, Uncertainty, and the Brain: The Science of Neuroeconomics* (MIT Press, Cambridge 2003).
 - [50] E. W. Piotrowski and J. Sladkowski, *The next stage: quantum game theory*, quant-ph/0308027.
 - [51] E. W. Piotrowski and J. Sladkowski, *Quantum computer: an appliance for playing market games*, quant-ph/0305017.
 - [52] G. Collum, *Systems of logical systems: Neuroscience and quantum logic*, Foundations of Science **7**, 49 (2002).